



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Journal Pre-proof

Clinical investigation of intestinal fatty acid-binding protein (I-FABP)
as a biomarker of SARS-CoV-2 infection



Rafael Simone Saia BPharm, PhD , Humberto Giusti BSc ,
Fábio Luís Silva MD, PhD ,
Karina de Jesus Bonicenha Pedroso RN ,
Maria Auxiliadora-Martins MD, PhD ,
Karen Mirna Loro Morejón MD, PhD ,
Augusto Marcussi Degiovani MD , Márcia Regina Cadelca MD ,
Anibal Basile-Filho MD, PhD

PII: S1201-9712(21)00761-X
DOI: <https://doi.org/10.1016/j.ijid.2021.09.051>
Reference: IJID 5734

To appear in: *International Journal of Infectious Diseases*

Received date: 4 August 2021
Revised date: 21 September 2021
Accepted date: 23 September 2021

Please cite this article as: Rafael Simone Saia BPharm, PhD , Humberto Giusti BSc , Fábio Luís Silva MD, PhD , Karina de Jesus Bonicenha Pedroso RN , Maria Auxiliadora-Martins MD, PhD , Karen Mirna Loro Morejón MD, PhD , Augusto Marcussi Degiovani MD , Márcia Regina Cadelca MD , Anibal Basile-Filho MD, PhD , Clinical investigation of intestinal fatty acid-binding protein (I-FABP) as a biomarker of SARS-CoV-2 infection, *International Journal of Infectious Diseases* (2021), doi: <https://doi.org/10.1016/j.ijid.2021.09.051>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2021 Published by Elsevier Ltd on behalf of International Society for Infectious Diseases.
This is an open access article under the CC BY-NC-ND license
(<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

HIGHLIGHTS

- SARS-CoV-2 induces long-lasting loss of enterocyte membrane integrity
- Elevated urinary I-FABP levels are associated with disease severity and poor outcome
- Damage to enterocytes is correlated to high IL-6 production

Journal Pre-proof

**Clinical investigation of intestinal fatty acid-binding protein (I-FABP) as a
biomarker of SARS-CoV-2 infection**

Rafael Simone Saia BPharm, PhD ^{1,*}, Humberto Giusti, BSc ¹, Fábio Luís Silva, MD, PhD ^{2,3},
Karina de Jesus Bonicenna Pedrosa, RN ⁴, Maria Auxiliadora-Martins, MD, PhD ², Karen Mirna
Loro Morejón, MD, PhD ⁴, Augusto Marcussi Degiovani, MD ⁵, Márcia Regina Cadelca, MD ⁶
and Anibal Basile-Filho, MD, PhD ²

¹ Laboratory of Intestinal Pathophysiology, Department of Physiology, Ribeirão Preto Medical
School, University of São Paulo, Ribeirão Preto, São Paulo, Brazil.

² Division of Intensive Care Medicine, Department of Surgery and Anatomy, Ribeirão Preto
Medical School, University of São Paulo, Ribeirão Preto, São Paulo, Brazil.

³ Intensive Care Unit, Fundação Hospital Santa Lydia, Ribeirão Preto, São Paulo, Brazil.

⁴ Hospital Unimed Ribeirão Preto, Ribeirão Preto, São Paulo, Brazil.

⁵ Intensive Care Unit, Santa Casa de Misericórdia, Ribeirão Preto, São Paulo, Brazil.

⁶ Intensive Care Unit, Hospital e Maternidade Santa Isabel, Jaboticabal, São Paulo, Brazil.

*** Corresponding Author and Reprint Requests:** Rafael Simone Saia, Department of
Physiology, Ribeirão Preto Medical School/USP. Avenida dos Bandeirantes, 3900. 14049-900 -
Ribeirão Preto, SP, Brazil. Fax: +55 16 3315-0221 Phone: +55 16 3315-3199.

E-mail: rssaia@fmrp.usp.br

Number of figures: 1

Number of tables: 1

Short Title: Damage to enterocytes in SARS-CoV-2

Disclosure Statement: The authors declare that there are no conflicts of interest that could be
perceived as prejudicing the impartiality of the research reported.

ABSTRACT

Objectives: SARS-CoV-2 exhibits tropism for the gastrointestinal tract, however lesions in enterocytes and their correlation with disease severity and patient prognosis are still unknown.

Methods: SARS-CoV-2 patients were enrolled in five medical centres in the São Paulo State, Brazil. Clinical characteristics and laboratory findings of patients were recorded. At admission, 7th and 14th day of hospitalisation, plasma and urine samples were collected. The levels of cytokines (IFN- γ , IL-6 and TNF- α) and the biomarker intestinal fatty acid-binding protein (I-FABP) were measured.

Results: COVID-19 patients displayed \approx 48-, 74- and 125-fold increased urinary I-FABP levels at admission ($n=283$; $p<0.001$), 7th ($n=142$; $p<0.01$) and 14th day ($n=75$; $p<0.01$) of hospitalisation, respectively. Critically ill patients and nonsurvivors showed higher I-FABP concentrations compared to those in the infirmary and survivors, respectively, within the same hospital stay periods. At admission, infected patients demonstrated enhanced production of plasma IFN- γ and IL-6. The ROC curve suggested I-FABP as a biomarker for disease severity of COVID-19 at admission ($p<0.0001$; Youden index=6.89; AUC=0.699). Patients with I-FABP \geq 6.89 showed higher IL-6 and C-reactive protein levels ($p<0.001$) at admission, as well as prolonged length of hospital stay.

Conclusions: Our findings revealed damage to enterocytes in SARS-CoV-2 infection, which is associated to illness severity, poor prognosis and exacerbated inflammatory response.

Keywords: intestinal epithelial cells, cytokines, inflammation, COVID-19, interleukin-

SHORT COMMUNICATION

Gastrointestinal (GI) symptoms including vomiting, abdominal discomfort and diarrhoea have been reported in coronavirus infections at the onset of hospitalisation, affecting 10-30% of patients (Parasa et al., 2020). SARS-CoV-2 RNA has been detected in faeces and stomach, duodenum and colon biopsies (Lin et al., 2020; Zhao et al., 2020). COVID-19 patients with ongoing diarrhoea have displayed enhanced concentration of faecal calprotectin (a marker of gut inflammation), as well as systemic interleukin (IL)-6 production, regardless of the viral load (Effenberger et al., 2020). Although SARS-CoV-2 exhibits tropism for the GI tract, it remains unknown whether lesions in the intestinal epithelium occur.

Intestinal fatty acid-binding protein (I-FABP) is a cytosolic protein expressed in mature enterocytes of the small and large intestines (Pelsers et al., 2003). A basal level of urinary I-FABP represents the physiological turnover rate of the epithelium, whilst its increment may indicate enterocytes cell damage. In sepsis and IBD, the increased urinary concentration of I-FABP has been confirmed as a reliable biomarker and predictor of disease reactivation, worse prognosis and clinical complications (Ho et al., 2020). Therefore, our study assessed the levels of I-FABP in hospitalised patients with SARS-CoV-2 infection and investigated its role as a predictor of disease severity and poor prognosis.

This prospective cohort study enrolled 283 inpatients from five medical centres (Hospital das Clínicas, Hospital Santa Lydia, Hospital Unimed and Santa Casa de Misericórdia in Ribeirão Preto city and also Hospital e Maternidade Santa Isabel in Jaboaticabal city), São Paulo State, Brazil (**Supplementary methods**). SARS-CoV-2 infection was confirmed through viral RNA detection in nasopharyngeal swabs by real-

time RT-PCR and the positive cases were recruited between April to September 2020. Comorbidities such as GI disorders (IBD, celiac disease and cancer), HIV, and alcohol and drugs addictions were exclusion criteria. At admission, 7th and 14th day of hospitalisation (at infirmary or intensive care unit), venous blood and urine samples were collected in the morning (0600-0800 AM), chilled and transferred to the research laboratory. Total blood and urine were centrifuged (1200 g, 15 min, 4 °C), the supernatants aliquoted and kept at -70°C for further analysis. Healthy volunteers were seronegative for the presence of anti-SARS-CoV-2 IgM and IgG antibodies.

I-FABP and cytokines (TNF- α , IFN- γ and IL-6) were quantified by ELISA, according to the manufacturer's instructions (R&D Systems, Minneapolis, MN, USA). Urinary data were normalised to creatinine concentrations (LabTest, Lagoa Santa, MG, Brazil). Continuous variables were expressed as median (interquartile range [IQR]) or 10-90 percentile range, and then compared using Mann-Whitney *U*-test or Kruskal-Wallis test (**Supplementary results**). Categorical variables were compared by Fisher's exact test or Pearson χ^2 test. Statistical significance was considered different when $p < 0.05$.

Table 1 lists the demographic, clinical characteristics, and laboratory findings of SARS-CoV-2 patients (survivors and nonsurvivors) and healthy volunteers. Nonsurvivors showed higher number of underlying comorbidities (cardiovascular and pulmonary diseases, diabetes, degenerative and malignancies), as well as deviations of the laboratory parameters (anaemia, leucocytosis, lymphocytopenia, D-dimers, CRP, LDH, serum creatinine and urea), requirement of mechanic ventilation and intensive care at admission. At 7th and 14th day of hospitalisation, aging, pre-existing diseases, leucocytosis and need for mechanic ventilation were relevant factors in nonsurvivor patients.

Noteworthy, COVID-19 patients displayed \approx 48-, 74- and 125-fold increased urinary I-FABP levels at admission, 7th and 14th day of hospitalisation, respectively (**Figure 1A**; $p<0.001$). Critically ill patients showed higher I-FABP concentrations compared to those in the infirmary within the same hospital stay periods (**Figure 1B**). Likewise, nonsurvivor patients were distinguished by increased I-FABP levels in contrast to survivors (**Figure 1C**). At admission, the inflammatory response was characterised by augmented plasma IFN- γ and IL-6 levels in SARS-CoV-2 patients (**Figure 1D**). Interestingly, urinary IL-6 concentrations were elevated in infected patients and nonsurvivors (**Figure 1E**). The ROC analysis indicated I-FABP as a good biomarker for disease severity, predictor for poor prognosis of COVID-19 at admission (**Figure 1F**; Youden index=6.89; $p<0.0001$) and prolonged hospital stay (**Figure 1G**). Patients with I-FABP \geq 6.89 ng/mg showed higher plasma and urine IL-6 and serum CRP concentrations, suggesting an interaction between inflammation and damage to enterocytes (**Figure 1G**).

Epithelial cells from ileum and colon express ACE2, the host cell entry receptor of coronaviruses, invading them for replication (Hoffmann et al., 2020) and shedding virions for several weeks after the onset of the illness (Zhao et al., 2020). Enteric symptoms reflect a break in the gut mucosa homeostasis, triggering the immune response and “cytokine release syndrome” (Effenberger et al., 2020). Soluble protein mediators, such as IFN- γ and IL-6, are key drivers of the inflammation-associated enterocyte damage and also responsible for increased gut mucosal permeability (Neurath, 2014; Wang et al., 2001). Indeed, our findings confirmed the detrimental contribution of SARS-CoV-2 to the loss of enterocyte membrane integrity, which is correlated with increased I-FABP levels (up to the 14th day after hospitalisation) and patient clinical condition. Moreover, patients whose I-FABP \geq 6.89 ng/mg also showed

exacerbated IL-6 production, which is a predictive cytokine of disease progression and ARDS complication (Santa Cruz et al., 2021). Our results revealed that increased I-FABP levels at admission may predict poor prognosis and SARS-CoV-2 illness severity, such as need for intensive care, mechanic ventilation and prolonged hospital stay.

Abbreviations list: ACE-2: angiotensin converting enzyme-2; ARDS: acute respiratory distress syndrome; AUC: area under curve; COVID-19: coronavirus disease of 2019; Cr: creatinine; CRP: C-reactive protein; ELISA: enzyme-linked immunosorbent assay; GI: gastrointestinal; HIV: human immunodeficiency virus; IBD: inflammatory bowel diseases; ICU: intensive care unit; IFN- γ : interferon- γ ; Ig: immunoglobulin; IL: interleukin; ROC curve: receiver operating characteristic curve; RT-PCR: reverse transcriptase polymerase chain reaction; SARS-CoV-2: severe acute respiratory syndrome coronavirus 2; TNF- α : tumor necrosis factor- α .

Funding: This study was supported by The Brazilian National Council for Scientific and Technological Development (CNPq, Grant #401337/2020-0), Ministry of Health and Ministry of Science, Technology and Innovation; The São Paulo Research Foundation (FAPESP, Grant #2020/10097-5); Foundation for Support to Education, Research and Assistance (FAEPA, Grant #416/2020). The manuscript content is solely the responsibility of the authors and does not necessarily represent the official view of the Research Foundations.

Ethics approval: The study protocol was approved by the Research Ethics Committee of Clinics Hospital of Ribeirão Preto Medical School, University of São Paulo (protocol CAAE 30794820.7.0000.5440) and all patients or family members gave the informed consent to participate in this research.

Patient and public involvement: Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Acknowledgments: We are grateful to the excellent support of the medical teams (doctors, nurses, physiotherapists and psychologists) in the hospitals involved in this study: 1) Santa Lydia Hospital: Dr. Marcelo César Carboneri, Dr. Walther de O. Campos Filho, Bárbara R. Bauduino Brandão, Eliane Bernardes de Queiróz, Luís F. de Melo Bougleux, Paula C. Trugillo Silva Neves, Renata C. Braulio da Silva, Silmara Miamoto, Sílvia Tolomeotti; 2) Unimed Ribeirão Preto Hospital: Dr. Gustavo Ribeiro de Oliveira, Dr. Plínio José E. de Camargo, Adriana de Fátima S. dos Santos, Bruna Maritan da Costa, Giovana A. Faggion Nomellini, Larissa Mil-Homens Albergaria; 3) Santa Casa de Misericórdia de Ribeirão Preto: Dr. Luiz C. Fontes Mega, Mariana C. Corrêa Marcolla, Rafael Cruz da Silva; and 4) Hospital e Maternidade Santa Isabel: Dr. Pedro Miguel Verardino, Edson Alves Julião, Nayara A. Poltronieri Correa.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Contributors: Study concept and design: RSS, HG, FLS, ABF. Involved in the patient care: FLS, MAM, KMLM, AMD, MRC. Patient recruitment: RSS, HG, FLS, KJBP, MAM, AMD, MRC. Laboratory Analyses: RSS, HG. Data collection: RSS, HG, FLS, KJBP. Statistical analysis: RSS, ABF. Drafting the manuscript: RSS. All the authors edited the manuscript and have approved the final version.

Preprint Server: The present results have not been published as a preprint version.

References

- Effenberger M, Grabherr F, Mayr L, Schwaerzler J, Nairz M, Seifert M, et al. Faecal calprotectin indicates intestinal inflammation in COVID-19. *Gut* 2020;69(8):1543-4. <https://10.1136/gutjnl-2020-321388>.
- Ho SS, Wall C, Gearry RB, Keenan J, Day AS. A Pilot Study Evaluating Novel Urinary Biomarkers for Crohn's Disease. *Inflam Intest Dis* 2020;5(4):212-20. <https://10.1159/000510682>.
- Hoffmann M, Kleine-Weber H, Schroeder S, Kruger N, Herrler T, Erichsen S, et al. SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is Blocked by a Clinically Proven Protease Inhibitor. *Cell* 2020;181(2):271-80 e8. <https://10.1016/j.cell.2020.02.052>.
- Lin L, Jiang X, Zhang Z, Huang S, Fang Z, Gu Z, et al. Gastrointestinal symptoms of 95 cases with SARS-CoV-2 infection. *Gut* 2020;69(6):997-1001. <https://10.1136/gutjnl-2020-321013>.
- Neurath MF. Cytokines in inflammatory bowel disease. *Nat Rev Immunol* 2014;14(5):329-42. <https://10.1038/nri3661>.
- Parasa S, Desai M, Thoguluva Chandrasekar V, Patel HK, Kennedy KF, Roesch T, et al. Prevalence of Gastrointestinal Symptoms and Fecal Viral Shedding in Patients With Coronavirus Disease 2019: A Systematic Review and Meta-analysis. *JAMA* 2020;3(6):e2011335. <https://10.1001/jamanetworkopen.2020.11335>.
- Pelsers MM, Namiot Z, Kisielewski W, Namiot A, Januszkiewicz M, Hermens WT, et al. Intestinal-type and liver-type fatty acid-binding protein in the intestine. Tissue distribution and clinical utility. *Clin Biochem* 2003;36(7):529-35. [https://10.1016/s0009-9120\(03\)00096-1](https://10.1016/s0009-9120(03)00096-1).
- Santa Cruz A, Mendes-Frias A, Oliveira AI, Dias L, Matos AR, Carvalho A, et al. Interleukin-6 Is a Biomarker for the Development of Fatal Severe Acute Respiratory Syndrome Coronavirus 2 Pneumonia. *Front Immunol* 2021;12:613422. <https://10.3389/fimmu.2021.613422>.

Wang Q, Fang CH, Hasselgren PO. Intestinal permeability is reduced and IL-10 levels are increased in septic IL-6 knockout mice. *Am J Physiol* 2001;281(3):R1013-23. <https://doi.org/10.1152/ajpregu.2001.281.3.R1013>.

Zhao F, Yang Y, Wang Z, Li L, Liu L, Liu Y. The Time Sequences of Respiratory and Rectal Viral Shedding in Patients With Coronavirus Disease 2019. *Gastroenterology* 2020;159(3):1158-60 e2. <https://doi.org/10.1053/j.gastro.2020.05.035>.

Journal Pre-proof




Figure 1. Urinary I-FABP as a prognostic biomarker of SARS-CoV-2 disease severity. (A) Urinary I-FABP concentration determined by ELISA in healthy volunteers and SARS-CoV-2 infected patients at admission, 7th and 14th day of hospitalisation (Kruskal-Wallis test and Dunn's multiple comparison post-test). (B) Urinary I-FABP concentration in SARS-CoV-2 patients in non-critical and critical clinical conditions at admission, 7th and 14th day of hospitalisation (Mann-Whitney test). (C) Urinary I-FABP concentration in SARS-CoV-2 survivor and nonsurvivor patients at admission, 7th and 14th day of hospitalisation (Mann-Whitney test). (D) Plasma cytokines levels of TNF- α , IFN- γ and IL-6 determined by ELISA at admission (Mann-Whitney test). (E) Urinary IL-6 concentration at admission in healthy volunteers, SARS-CoV-2 infected patients, survivors and nonsurvivors (Mann-Whitney test). (F) Comparison of ROC analysis of urinary I-FABP (AUC=0.699, 95% CI=0.636-0.762, $p<0.0001$), urinary IL-6 (AUC=0.573, 95% CI=0.476-0.671, $p=0.14$), plasma IL-6 (AUC=0.509, 95% CI=0.436-0.582, $p=0.82$), plasma IFN- γ (AUC=0.523, 95% CI=0.445-0.602, $p=0.54$) and serum CRP (AUC=0.591, 95% CI=0.519-0.662, $p=0.01$) for SARS-CoV-2 survivor and non-survivor patients. Cr: creatinine. (G) Plasma and urinary IL-6 concentrations, serum CRP levels at admission and length of hospital stay in SARS-CoV-2 infected

patients divided according to the Youden index=6.89 ng/mg creatinine (low I-FABP<6.89 and high I-FABP \geq 6.89 ng/mg) obtained from the ROC curve in (F) (Mann-Whitney test). Results shown as bars and violin plots are expressed as median and 10-90 interquartile range. *n* represents the number of patients per group. *P* values indicate the statistical significance between groups.

Journal Pre-proof

Table 1. Demographic, clinical characteristics and laboratory parameters of healthy volunteers and SARS-CoV-2 infected patients (Survivors and Nonsurvivors)

Parameters	Healthy Volunteers (n=37)	SARS-CoV-2 infected patients					
		Admission (n=283)		7 th day (n=142)		14 th day (n=75)	
		Survivors (n=181)	Nonsurvivors (n=102)	Survivors (n=76)	Nonsurvivors (n=66)	Survivors (n=42)	Nonsurvivors (n=33)
Gender M/F (%)	33/67 ^γ	60/40	52/48	57/43	51/49	67/33	48/52 ^b
Age, years ^{&}	55 [49-60] ^α	61 [50-71]	69 [62-77] ^c	62 [54-70]	68 [58-77] ^b	61 [53-70]	68 [58-77] ^b
Ethnicity (C/AA/H) (%)	87/8/5 ^β	70/15/15	80/16/4 ^a	75/18/7	86/2/12 ^c	74/12/14	85/1/14 ^b
BMI, kg/m ² ^{&}	27.5 [24.5-30.3] ^α	30.5 [26.7-34.2]	28.7 [26.0-35.3]	31.5 [27.5-35.2]	29.0 [25.9-35.8]	29.8 [26.8-34.4]	28.9 [25.5-37.6]
MAP, mmHg ^{&}	90.0 [80.0-96.7]	93.3 [80.5-99.1]	88.0 [76.3-96.0]	91.0 [78.3-98.3]	87.0 [75.3-96.7]	91.0 [78.3-98.3]	88.5 [77.9-104.2]
HR, beats/min ^{&}	75 [69-79] ^γ	85 [73-92]	86 [77-98]	85 [76-96]	90 [79-98]	85 [76-96]	91 [83-104] ^a
Underlying Comorbidities, no. (%)							
Smoker/ex-smoker	0 (0) ^γ	29 (16.0)	31 (30.4) ^b	15 (19.7)	16 (24.2)	8 (19.0)	8 (24.2)
Arterial hypertension	10 (27.0) ^γ	99 (54.7)	74 (72.5) ^b	51 (67.1)	50 (75.6)	28 (66.7)	30 (90.9) ^b
Cardiovascular diseases	1 (2.7) ^γ	32 (17.7)	31 (30.4) ^a	18 (23.7)	22 (33.3)	11 (26.2)	10 (30.3)
Diabetes	2 (5.4) ^γ	61 (33.7)	50 (49.0) ^a	33 (43.4)	29 (43.9)	19 (45.2)	18 (54.5)
Pulmonary diseases	0 (0) ^β	9 (5.0)	15 (14.7) ^b	2 (2.6)	7 (10.6)	2 (4.8)	4 (12.1)
Chronic kidney diseases	0 (0)	7 (3.9)	9 (8.8)	2 (2.6)	5 (7.6)	0 (0)	4 (12.1) ^a
Thyroid disease	1 (2.7)	11 (6.1)	12 (11.8)	9 (11.8)	9 (13.6)	4 (9.5)	4 (12.1)
Dyslipidaemia	2 (5.4)	23 (12.7)	16 (15.7)	15 (19.7)	11 (16.7)	9 (21.4)	4 (12.1)
Degenerative disorders	0 (0) ^γ	1 (0.5)	9 (8.8) ^c	0 (0)	4 (6.1) ^a	0 (0)	0 (0)
Neurological diseases	1 (2.7)	12 (6.6)	6 (5.9)	8 (10.5)	7 (10.6)	5 (11.9)	4 (12.1)
Malignancies	0 (0) ^γ	2 (1.1)	15 (14.7) ^c	1 (1.3)	9 (13.6) ^b	1 (2.4)	6 (18.2) ^a
Other diseases	3 (8.1) ^α	8 (3.9)	13 (12.7) ^a	5 (6.6)	8 (12.1)	3 (7.1)	5 (15.2)

Haematological Indexes &

Haemoglobin, g/dL	14.8 [14.0-15.2] ^γ	12.6 [11.3-14.0]	11.6 [10.0-13.3] ^b	11.7 [10.2-12.9]	10.0 [8.5-12.3] ^c	9.7 [8.7-11.3]	9.2 [8.2-11]
Erythrocytes, 10 ⁶ /μL	4.5 [4.42-4.90] ^a	4.3 [3.9-4.7]	4.2 [3.6-4.6]	4.0 [3.6-4.4]	3.67 [3.0-4.3] ^b	3.3 [2.9-3.8]	3.12 [2.74-3.69]
Leukocytes, 10 ³ /μL	5.38 [4.35-5.71] ^γ	8.2 [6.1-11.08]	11.4 [7.78-15.2] ^c	11.45 [8.23-16.03]	14.5 [10.2-23.2] ^b	10.2 [8.2-12.8]	12.6 [9.34-19.83] ^a
Neutrophils, 10 ³ /μL	2.65 [1.97-3.08] ^γ	6.01 [4.2-8.7]	8.89 [6.01-12.9] ^c	9.00 [6.0-12.4]	11.8 [7.45-19.85] ^c	7.8 [4.3-9.8]	10.75 [6.31-17.05] ^a
Monocytes, 10 ³ /μL	0.34 [0.27-0.38]	0.36 [0.22-0.54]	0.4 [0.27-0.68]	0.5 [0.3-0.7]	0.5 [0.31-0.8]	0.5 [0.3-0.8]	0.5 [0.3-0.8]
Lymphocytes, 10 ³ /μL	1.94 [1.47-2.17] ^β	1.1 [0.7-1.44]	0.89 [0.6-1.34] ^a	1.23 [0.71-1.79]	1.0 [0.6-1.59]	1.51 [1.0-2.1]	1.27 [0.93-1.82]
Platelets, 10 ³ /μL	209 [181-247]	230 [173-304]	230 [181-287]	271 [224-358]	250 [172-322] ^a	232 [165-304]	235 [143-327]
D-dimer, μg/mL [†]	< 0.5 ^γ	1.4 [0.59-3.25]	3.17 [1.84-6.84] ^c	2.31 [1.32-3.32]	2.61 [1.93-3.58]	2.51 [1.74-4.37]	3.4 [2.55-10.11] ^a

Biochemical Indexes &

Glycaemia, mg/dL	78 [75-87] ^γ	155 [119-229]	168 [138-230]	170 [116-234]	176 [127-232]	142 [117-198]	133 [114-208]
CRP, mg/L	0.3 [0.2-0.4] ^γ	41.7 [16.7-109.3]	76.8 [20.1-163.3] ^a	21.3 [7.0-54.0]	33.6 [9.1-109.7]	67.0 [21.1-143.5]	53.5 [12.1-180.0]
ALT, U/L	20.4 [13.8-28.6] ^γ	45.5 [27.3-66.3]	39.5 [22.5-67.2]	61.0 [43.0-95.7]	58 [31.0-110.7]	66.0 [43.1-106.5]	57.1 [45.3-101.7]
AST, U/L	21.0 [19.0-27.0] ^γ	42.0 [30.0-65.9]	50.0 [29.0-71.2]	42.3 [31.0-55.3]	47.1 [33.0-86.0]	42.5 [29.0-69.5]	39.2 [34.9-111.5]
LDH, U/L	183.0 [172-189] ^γ	340.0 [263-466.4]	437.1 [293.6-675] ^b	345.0 [279-470]	408.3 [327.9-607] ^b	326.0 [259-401]	392.2 [285-571.9]
Serum creatinine, mg/dL	0.8 [0.7-0.9] ^a	1.0 [0.8-1.3]	1.43 [0.97-2.63] ^c	1.0 [0.8-1.5]	1.56 [1.01-2.87] ^c	1.1 [0.8-1.5]	1.66 [0.87-2.12] ^a
Serum urea, mg/dL	31.0 [28.0-38.0] ^a	45.0 [31.2-62.0]	74.9 [46.6-119.1] ^c	62.0 [40.0-99.9]	111.9 [56.2-167.8] ^c	62.0 [41.0-108.2]	102.6 [46.3-155.3] ^a

Respiratory Indexes &

pH [†]	7.4 [7.35-7.45]	7.4 [7.32-7.44]	7.32 [7.23-7.40] ^c	7.4 [7.34-7.43]	7.34 [7.25-7.4]	7.4 [7.35-7.44]	7.33 [7.28-7.4]
PaO ₂ , mmHg [†]	90.0 [80-100]	83.9 [71.0-95.2]	78.4 [67.7-91.5]	72.5 [67.0-82.2]	75.4 [68.9-83.3]	73.4 [64.6-80.7]	73.0 [67.4-80.4]
PaCO ₂ , mmHg [†]	40.0 [35.0-45.0]	34.8 [27.8-45.0]	42.2 [34.0-48.6] ^b	44.1 [38.9-51.9]	45.4 [35.0-51.8]	38.4 [33.4-46.1]	42.5 [32.6-49.2]
SaO ₂ , % [†]	96.5 [95.0-98.0]	96.4 [93.8-97.1]	95.6 [92.3-97.1] ^a	96.4 [94.1-97.2]	95.7 [94.1-97.2]	96.0 [94.8-97.0]	95.5 [93.0-97.0]
Lactate, mmol/L [†]	1.25 [0.5-2.0]	2.0 [1.45-2.6]	2.0 [1.42-2.75]	1.89 [1.37-2.6]	1.77 [1.28-2.6]	1.80 [1.4-2.4]	1.51 [0.76-2.71]

Treatments, no. (%)

Antibiotic therapy	-	132 (72.9)	102 (100.0) ^c	59 (77.6)	51 (77.3)	40 (95.2)	33 (100.0)
Antiviral therapy	-	25 (13.8)	12 (11.8)	2 (2.6)	1 (1.5)	0 (0)	0 (0)
Corticosteroids	-	123 (68.0)	81 (79.4) ^a	58 (76.3)	51 (77.3)	21 (50.0)	24 (72.7)
Anticoagulants	-	112 (61.9)	93 (91.2) ^c	75 (98.7)	55 (83.3) ^c	41 (97.6)	33 (100.0)
Hydroxychloroquine	-	12 (6.6)	4 (3.9)	3 (3.9)	3 (4.5)	0 (0)	0 (0)

Hospitalisation
Characteristics

Length of hospital stay (days)	-	11 [4-22]	13 [8-21] ^a	-	-	-	-
ICU length of stay (days)	-	0 [0-12]	11 [6-19] ^c	-	-	-	-
Severe/critical care type, no. (%) [†]	-	73 (40.3)	88 (86.3) ^c	54 (71.1)	63 (95.5) ^c	35 (83.3)	33 (100.0) ^a
Mechanic ventilation, no. (%)	-	60 (33.1)	73 (71.6) ^c	52 (68.4)	59 (89.4) ^b	28 (66.7)	33 (100.0) ^c

Legend: ALT (alanine aminotransferase); AST (aspartate aminotransferase); BMI (body mass index); CRP (C-reactive protein); Ethnicity (C = Caucasian, AA = Afro American, H = Hispanic); HR (heart rate); ICU (intensive care unit); LDH (lactate dehydrogenase); MAP (mean arterial pressure); PaCO₂ (partial pressure of carbon dioxide in arterial blood); PaO₂ (partial pressure of oxygen in arterial blood); SaO₂ (arterial oxygen saturation).

[&] Values expressed as median [interquartile range]

[†] Healthy volunteers with standardized reference values

[#] Severe/critical care type was defined as respiratory failure requiring mechanical ventilation and ICU admission

^a $p < 0.05$, ^b $p < 0.01$ and ^c $p < 0.001$ vs. SARS-CoV-2 patients (survivors and nonsurvivors) at the admission

^a $p < 0.05$, ^b $p < 0.01$ and ^c $p < 0.001$ vs. survivors SARS-CoV-2 patients at the respective period of hospitalisation